

# Measurement of urea synthesis in cultured stem cell-derived hepatocyte-like cells

Created on: 08-03-2019 - Last modified on: 28-02-2022

## Organisation

**Name of the organisation** Vrije Universiteit Brussel (VUB)

**Department** Pharmaceutical and Pharmacological Sciences

**Specific Research Group or Service** In Vitro Toxicology and Dermato-Cosmetology

**Country** Belgium

**Geographical Area** Brussels Region

## SCOPE OF THE METHOD

<b>The Method relates to</b>	Human health
<b>The Method is situated in</b>	Basic Research
<b>Type of method</b>	In vitro - Ex vivo
<b>Specify the type of cells/tissues/organs</b>	Human stem cells

## DESCRIPTION

### Method keywords

Stem cells

Hepatocytes

Hepatotoxicity

urea

### Scientific area keywords

hepaticdifferentiation

hepatic toxicity  
Hepatotoxicity  
Cell culture  
cellular programming

### **Method description**

The present standard procedure describes a protocol for measuring the urea concentration in supernatant of human stem cell-derived hepatocyte-like cells. This procedure relies on a chromogenic reagent that forms a colored complex specifically with urea. The latter can be measured and is directly proportional to the urea concentration in the sample.

### **Lab equipment**

Biosafety cabinet;  
Multiplate reader;  
Thermostated bath.

### **Method status**

History of use  
Internally validated

## **PROS, CONS & FUTURE POTENTIAL**

### **Advantages**

The current protocol represents a simple and direct method to quantitatively measure the urea concentration in human stem cell-derived hepatocyte-like cell cultures. This assay has no harmful effect on the cultured cells. Therefore, after incubation of the cells with the substrate (ammonium chloride (NH<sub>4</sub>Cl)), the cultures can be maintained.

### **Future & Other applications**

Can be applied to other types of *in vitro* systems of hepatocytes.

## **REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION**

## References

- Gómez-Lechón MJ, Lahoz A, Gombau L, Castell JV, Donato MT. (2010) In vitro evaluation of potential hepatotoxicity induced by drugs. *Curr Pharm Des* 16:1963-77
- Lavon N, Benvenisty N. (2005) Study of hepatocyte differentiation using embryonic stem cells. *J Cell Biochem* 96:1193-202
- Roelandt P, Pauwelyn KA, Sancho-Bru P, Subramanian K, Ordovas L, Vanuytsel K, Geraerts M, Firpo M, De Vos R, Fevery J, Nevens F, Hu WS, Verfaillie CM. (2010) Human Embryonic and Rat Adult Stem Cells with Primitive Endoderm-Like Phenotype Can Be Fated to Definitive Endoderm, and Finally Hepatocyte-Like Cells. *PLoS One*; 5(8): e12101
- Snykers S, Vanhaecke T, De Becker A, Papeleu P, Vinken M, Van Riet I, Rogiers V. (2007) Chromatin remodeling agent trichostatin A: a key-factor in the hepatic differentiation of human mesenchymal stem cells derived of adult bone marrow. *BMC Dev Biol* 7:24
- Snykers S, De Kock J, Rogiers V, Vanhaecke T. (2009) In vitro differentiation of embryonic and adult stem cells into hepatocytes: state of the art. *Stem Cells* 27:577-605

## Associated documents

[Urea synthesis.doc](#)

*Coordinated by*



*Financed by*

