

# Lentiviral reprogramming of human umbilical cord-derived mesenchymal stromal cells towards hepatocyte-like cells

Created on: 20-03-2019 - Last modified on: 28-02-2022

#### **Contact person**

Karolien Buyl

#### Organisation

Name of the organisation Vrije Universiteit Brussel (VUB)
Department Pharmaceutical and Pharmacological Sciences
Specific Research Group or Service In Vitro Toxicology and Dermato-Cosmetology
Country Belgium
Geographical Area Brussels Region

# SCOPE OF THE METHOD

The Method relates to	Human health
The Method is situated in	Basic Research
Type of method	In vitro - Ex vivo
Specify the type of cells/tissues/organs	umbilical cord

## DESCRIPTION

#### Method keywords

human adult stem cells hepatocyte-like cells lentiviral reprogramming

#### umbilical cord

#### Scientific area keywords

in vitro liver model Drug-induced liver injury (DILI) human hepatocyte-like cells liver enriched transcription factor human adult stem cells

#### Method description

Human umbilical cord-derived mesenchymal stromal cells (hUC-MSCs) express several key liver-specific transcription factors as well as hepatic progenitor markers. However, they still lack the hepatocyte nuclear factors 1-alpha (HNF1a) and 4-alpha (HNF4a), indispensable for their reprogramming towards hepatocyte-like cells. This method comprises the reprogramming of hUC-MSCs towards hepatocyte-like cells through HNF1a lentiviral over-expression. Whole genome microarray analysis revealed that the expression of the nuclear receptor retinoid X receptor (RXR) gamma and the nuclear transcription factor HNF4a, in HNF1a-transduced hUC-MSCs, was significantly upregulated compared to the control conditions. This expression was even higher than found in human hepatocytes. The same was observed for the uridine 5'-diphospho-glucuronosyltransferase (UGT) 1A family. Further, a significant upregulation was observed for alpha-foetoprotein (AFP), alpha1-antitrypsin (A1-AT), the phase I biotransformation enzymes cytochrome P450 (CYP) 1A2 and CYP2A6 and the drug transporter multidrug resistance protein (MDR) 1.

#### Lab equipment

Incubator (37 ± 1°C, 90 ± 5% humidity, 5.0 ± 1% CO2/air); Type 2 laminar airflow HEK293T cells; Ultracentrifuge Water bath (37 ± 1°C); Inverse-phase contrast microscope; Pipettes; Pipettors; Colorimetric reverse transcriptase assay; Human hepatocytes; Affymetrix microarray technologies; Partek Genomics Suite Software.

## Method status

Still in development

# **PROS, CONS & FUTURE POTENTIAL**

## Advantages

Persistent expression of HNF1a transcription factor in hUC-MSCs; Endogenous induction of HNF4a expression.

# Challenges

Genomic integration of the lentiviral vector.

# Modifications

Usage of a non-integrative reprogramming method e.g. mRNA transfection.

# Future & Other applications

Generation of hepatocyte-like cells for the development of functional human liverbased *in vitro* models for pharmaco-toxicological purposes.

# **REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION**

Coordinated by







Financed by



