

Artificial cells with *Xenopus laevis* eggs cell-free extracts

Created on: 08-07-2025 - Last modified on: 22-07-2025

Contact person

Martina Boiardi

Organisation

Name of the organisation Katholieke Universiteit Leuven (KUL)

Department Cellular and Molecular Medicine

Specific Research Group or Service Laboratory of Dynamics in Biological Systems

Country Belgium

Geographical Area Flemish Region

Name of the organisation Cellular and Molecular Medicine - KU Leuven

Department Cellular and Molecular Medicine

Specific Research Group or Service Laboratory of Dynamics in Biological Systems

Country Belgium

SCOPE OF THE METHOD

The Method relates to	Other: Cell cycle fundamental research
The Method is situated in	Basic Research
Type of method	In vitro - Ex vivo
Species from which cells/tissues/organs are derived	<i>Xenopus laevis</i> frog
Type of cells/tissues/organs	eggs

DESCRIPTION

Method keywords

Microfluidic devices
droplets
eggs
cell free extracts
microemulsion
ultracentrifuge
fluorescence microscope
timelapse

Scientific area keywords

cell biology
cell cycle
Droplet-based microfluidics
cytoskeleton

Method description

The large (1mm diameter), easily accessible eggs of the frog *Xenopus laevis* (100-1000 eggs at once per frog) offer the opportunity to reconstitute cell cycle events *in-vitro* by generating cell-free extracts which retain all the biochemical components that regulate cell cycle progression, as well as all the organelles and cytoskeletal networks. In summary, ovulation is induced in frogs by subcutaneous injection of chorionic gonadotropin. After about 16 hours, eggs are collected, inspected for quality, washed, and processed using several centrifugation steps to obtain the cytoplasmic cell-free extract. Extracts can either be arrested in a cell-cycle phase, or being cycling (i.e., oscillating between interphase and mitosis). Cycling extracts are obtained by activating the eggs with calcium ionophore, which mimics fertilization and activates the biochemical processes of the cell cycle. To mimic cellular behavior, experiments require cell-sized compartments with realistic shapes and boundaries. This is achieved in two distinct ways: (i) by encapsulating extracts in surfactant-stabilized droplets, termed artificial cells, formed by vortexing frog egg extracts with surfactants and oil; (ii) by generating droplets using droplets-microfluidics. Cell cycle event are observed and recorded using timelapse fluorescence microscopy.

Lab equipment

Cell-free extract preparation:

- ultracentrifuge

Droplets generation:

- vortex or
- microfluidic chips for droplets production
- microfluidic pumps
- microscope with high-speed camera for tracking droplets production

Imaging:

- fluorescence microscope for timelapse
- multichannel
- multipoint microscopy

Method status

History of use

Published in peer reviewed journal

PROS, CONS & FUTURE POTENTIAL

Advantages

- Frogs are injected with hormones once every 3 months with a minimally invasive procedure,
- Frogs can be kept in the lab for 5 years (or longer if quality of the eggs is not compromised),
- 100-1000 eggs per frog allows to obtain up to 2 mL of cell-free extract,

- Cell cycle events in cycling cell-free extracts are fast. A cell cycle lasts about 30min/1h (this is because in the early embryo cellular cleavages have a period of 30 min). When imaging for 18h, several cell cycle events can be observed,
- This method is well established since the '80s, and there is a lot of literature.

Challenges

- Apoptotic eggs must be removed to avoid compromising the experiment and obtaining a non-functional extract,
- During heatwaves it is recommended to work in temperature controlled rooms, because the cell-free extract quickly becomes apoptotic above 25 degrees,
- During heatwaves the quality of eggs may drop, as well as their quantity, even though the temperature at which frogs are housed is kept under control all year long.

Future & Other applications

Cell free extracts are not only used for studying cell cycle regulation. They have been used also for studying cytoskeletal structures (e.g., mitotic spindles).

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

- (1) Andrew W. Murray. Cell Cycle Extracts. *Methods in Cell Biology*, 36:581–605, 1991. ISSN 0091679X. doi: 10.1016/S0091-679X(08)60298-8
- (2) Oshri Afanзар, Garrison K Buss, Tim Stearns, and Jr Ferrell, James E. The nucleus serves as the pacemaker for the cell cycle. *eLife*, 9:e59989, dec 2020. ISSN 2050-084X
- (3) Felix E Nolet, Alexandra Vandervelde, Arno Vanderbeke, Liliana Piñeros, Jeremy B Chang, and Lendert Gelens. Nuclei determine the spatial origin of mitotic waves. *eLife*, 9:e52868, may 2020. ISSN 2050-084X.
- (4) Ye Guan, Zhengda Li, Shiyuan Wang, Patrick M Barnes, Xuwen Liu, Haotian Xu, Minjun Jin, Allen P Liu, and Qiong Yang. A robust and tunable mitotic oscillator in artificial cells. *eLife*, 7:e33549, apr 2018. ISSN 2050-084X.
- (5) Gembu Maryu and Qiong Yang. Nuclear-cytoplasmic compartmentalization of cyclin b1-cdk1 promotes robust timing of mitotic events. *Cell Reports*, 41(13):111870, 2022. ISSN 2211-1247.

Coordinated by



Financed by

