

# Measurement of the extracellular release of adenosine triphosphate in cultured primary rat hepatocytes

*Commonly used acronym: ATP measurement*

*Created on: 12-03-2019 - Last modified on: 08-11-2019*

## SCOPE OF THE METHOD

<b>The Method relates to</b>	Animal health
<b>The Method is situated in</b>	Basic Research
<b>Type of method</b>	In vitro - Ex vivo
<b>This method makes use of</b>	Animal derived cells / tissues / organs
<b>Species from which cells/tissues/organs are derived</b>	rat
<b>Type of cells/tissues/organs</b>	primary hepatocytes

## DESCRIPTION

### Method keywords

ATP

cytotoxicity

bioluminescent determination

extracellular ATP

### Scientific area keywords

Hepatotoxicity

liver

cholestasis

Steatosis

### **Method description**

ATP transports chemical energy within cells by serving as a substrate for kinases and as such fulfills a vital function in numerous cellular processes such as cell injury and subsequent cell death. ATP is therefore a crucial player in these events that are results of intracellular stress. Hepatotoxic chemical compounds can cause intracellular stress. The general cytotoxicity of compounds can be estimated through the bioluminescent assessment of extracellular release of ATP. As such, this procedure relies on two reactions in which firefly luciferase catalyzes the oxidation of luciferin to oxyluciferin and whereby ATP is consumed and light becomes emitted. The latter can be measured and is proportional to the amount of ATP present outside cells.

### **Lab equipment**

Multiplate reader (Victor, 1420 Multilabel counter, PerkinElmer, Belgium)

### **Method status**

History of use

### **PROS, CONS & FUTURE POTENTIAL**

#### **Advantages**

Easy to apply method to quantitatively characterize extracellular ATP release and hence cell injury in primary hepatocyte cultures.

#### **Challenges**

Increased extracellular levels of ATP do not specifically indicate cell death by either apoptosis or necrosis. Ideally this method should be combined with established tests, such as the monitoring of cell proliferation potential and mitochondrial function, which can be done by measurement of the incorporation of 5-bromo-2'-deoxyuridine

(BrdU) during DNA synthesis and by addressing an 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, respectively.

## **REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION**

### **References**

Eguchi Y., Shimizu S. and Tsujimoto Y. (1997) Intracellular ATP levels determine cell death fate by apoptosis or necrosis. *Cancer Research* 57:1835-1840

Elaut G., Henkens T., Papeleu P., Snykers S., Vinken M., Vanhaecke T. and Rogiers V. (2006) Molecular mechanisms underlying the dedifferentiation process of isolated hepatocytes and their cultures. *Current Drug Metabolism* 7: 629-660

Golstein P. and Kroemer G. (2007) Cell death by necrosis: towards a molecular definition. *Trends in Biochemical Sciences* 32: 37-43

Gómez-Lechón M.J., Lahoz A., Gombau L., Castell J.V. and Donato M.T. (2010) In vitro evaluation of potential hepatotoxicity induced by drugs. *Current Pharmaceutical Design* 16: 1963-1977

Jaeschke H., Gores G.J., Cederbaum A.I., Hinson J.A. Pessayre D. and Lemasters J.J. (2002) Mechanisms of hepatotoxicity. *Toxicological Sciences* 65: 166-176. McKim J.M. (2010) Building a tiered approach to in vitro predictive toxicity screening: a focus on assays with in vivo relevance. *Combinatorial Chemistry and High Throughput Screening* 13: 188-206

Papeleu P., Vanhaecke T., Henkens T., Elaut G., Vinken M., Snykers S. and Rogiers V. (2006) Isolation of rat hepatocytes. *Methods in Molecular Biology* 320: 229-237

Seglen P.O. (1976) Preparation of isolated rat liver cells. *Methods in Cell Biology* 13: 29-83

Tsujimoto Y. (1997) Apoptosis and necrosis: intracellular ATP level as a determinant for cell death modes. *Cell Death and Differentiation* 4: 429-434

Van Calsteren V. (2010) Analyse van de herhaalde toedieningstoxiciteitstesten met cosmetische ingrediënten bestudeerd op Europees niveau. Dissertation Master in Pharmaceutical Sciences - Vrije Universiteit Brussel, Rogiers V. (promotor) and Pauwels M. (co-promotor)

Vinken M., Decrock E., De Vuyst E., Leybaert L., Vanhaecke T. and Rogiers V. (2009)  
Biochemical characterisation of an in vitro model of hepatocellular apoptotic cell  
death. Alternatives to Laboratory Animals 37: 209-218

### Associated documents

## PARTNERS AND COLLABORATIONS

### Organisation

**Name of the organisation** Vrije Universiteit Brussel

**Department** Pharmaceutical and Pharmacological Sciences (FARM)

**Specific Research Group or Service** In Vitro Toxicology and Dermato-cosmetology

**Country** Belgium

**Geographical Area** Brussels Region

*Coordinated by*



*Financed by*

