

Establishment of sandwich cultures of primary human hepatocytes

Created on: 14-03-2019 - Last modified on: 08-11-2019

SCOPE OF THE METHOD

The Method relates to	Human health
The Method is situated in	Basic Research
Type of method	In vitro - Ex vivo
This method makes use of	Human derived cells / tissues / organs
Specify the type of cells/tissues/organs	Primary human hepatocytes

DESCRIPTION

Method keywords

Sandwich cultures of hepatocytes

Scientific area keywords

Drug-induced cholestasis

Method description

This method describes a well-known optimised human *in vitro* model of drug-induced cholestasis. Cryopreserved primary human hepatocytes are cultured between two layers of extracellular matrix scaffold, which will delay dedifferentiation and allows to restore cell-extracellular matrix interactions. The sandwich culture method can be

applied to both single cell culture dishes and multi-well plates, thus providing an opportune model for high-throughput screening.

Lab equipment

Method status

Still in development

PROS, CONS & FUTURE POTENTIAL

Advantages

Suitable for long-term exposure ;

Restored cell polarity ;

Presence of cell-ECM interactions ;

Formation of functional bile canalicular network ;

Maintain functional expression levels of transport proteins and xenobiotic metabolism enzymes ;

Applicable for quantifying and detecting cholestatic liabilities.

Challenges

Mass transfer barrier ;

Difficult to culture in 96-well plates ;

Require daily medium renewal due to accumulating toxic metabolites ;

Hypoxic environment.

Modifications

The model is already modified by introducing a renewal of the collagen layer every 3-4 days. As a result, the model shows an extended cultivation regime up to 14 days (Parmentier et al. 2013).

Future & Other applications

The model could be used to assess the overall hepatotoxic potential of drugs, cosmetics, biocides or food additives.

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

Gijbels E., Vilas-Boas V., Deferm N. et al. (2019) Mechanisms and in vitro models of drug-induced cholestasis. Archives of Toxicology (submitted)

Gijbels E., Vanhaecke T., Vinken M. (2019) Establishment of sandwich cultures of primary human hepatocytes. Methods in Molecular Biology - Protocols in Experimental Cholestasis Research (accepted)

Other references you can find in attached document

Associated documents

[Manuscript.docx](#)

PARTNERS AND COLLABORATIONS

Organisation

Name of the organisation Vrije Universiteit Brussel

Department Pharmaceutical and Pharmacological Sciences (FARM)

Specific Research Group or Service In Vitro Toxicology and Dermato-cosmetology

Country Belgium

Geographical Area Brussels Region

Coordinated by



Financed by

