

Isolation and cultivation of adipose tissuederived mesenchymal stromal cells

Commonly used acronym: AT-MSC

Created on: 20-03-2019 - Last modified on: 28-02-2022

Contact person

Joery De Kock

Organisation

Name of the organisation Vrije Universiteit Brussel (VUB)

Department Pharmaceutical and Pharmacological Sciences

Specific Research Group or Service In Vitro Toxicology and Dermato-Cosmetology

Country Belgium

Geographical Area Brussels Region

SCOPE OF THE METHOD

| The Method relates to | Animal health |
|--|--|
| The Method is situated in | Basic Research |
| Type of method | In vitro - Ex vivo |
| Specify the type of cells/tissues/organs | adipose tissue-derived mesenchymal stromal cells |

DESCRIPTION

Method keywords

adipose tissue

Stem cells

mesenchymal stromal cells

isolation

cultivation

Scientific area keywords

stem cell culture

stem cell isolation

mesenchymal stromal cells

Method description

Approximately 125 g of processed adipose tissue is incubated for 90 minutes at 37°C in dissociation medium (1:1) consisting of 1% (v/v) bovine serum albumin and 1 mg/mL collagenase A in phosphate buffered saline (PBS). After two filtration steps, the filtrate is carefully brought on top of 15 mL of Histopaque®-1077. Upon centrifugation for 20 minutes at 1000 g (4°C), the top layer is removed and the AT-MSC are collected in 50 mL PBS/BSA (1%). This procedure is carried out separately on two pieces of adipose tissue. Typically 5 - 20 x 10E7 viable cells are obtained per 250 g of processed adipose tissue. The isolated AT-MSC are then (sub)cultured as a monolayer in AT-MSC growth medium for 2 weeks, consisting of Dulbecco's Modified Eagle Medium supplemented with 10% (v/v) foetal bovine serum (FBS), 50 μg/mL streptomycin sulphate, 7.33 IU/mL benzyl penicillin and 2.5 μg/mL fungizone. Cell cultures are incubated at 37°C in a 5% (v/v) CO2 humidified atmosphere and passaged at subconfluency using TrypLE® express. Growth media is changed every 3 days.

Lab equipment

Biosafety cabinet level 2;

Cell incubator:

Centrifuge.

Method status

History of use

Internally validated

Published in peer reviewed journal

PROS, CONS & FUTURE POTENTIAL

Advantages

Robust isolation method for adipose tissue-derived mesenchymal stromal cells.

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

De Kock J, Najar M, Bolleyn J, Al Battah F, Rodrigues RM, Buyl K, Raicevic G, Govaere O, Branson S, Meganathan K, Gaspar JA, Roskams T, Sachinidis A, Lagneaux L, Vanhaecke T, Rogiers V. (2012) Mesoderm-derived stem cells: the link between the transcriptome and their differentiation potential. Stem Cells Dev. 21(18):3309-23

Najar M, Rodrigues RM, Buyl K, Branson S, Vanhaecke T, Lagneaux L, Rogiers V, De Kock J. (2014) Proliferative and phenotypical characteristics of human adipose tissue-derived stem cells: comparison of Ficoll gradient centrifugation and red blood cell lysis buffer treatment purification methods. Cytotherapy. 16(9):1220-8











