

# Transdermal and transmucosal kinetics using FDC

Created on: 10-10-2019 - Last modified on: 08-11-2019

## **Contact person**

Yorick Janssens

# **Organisation**

Name of the organisation Ghent University (UGent)

**Department** Pharmaceutical analysis

**Country** Belgium

Geographical Area Flemish Region

#### **SCOPE OF THE METHOD**

The Method relates to	Human health
The Method is situated in	Basic Research
Type of method	In vitro - Ex vivo
Specify the type of cells/tissues/organs	Skin

## **DESCRIPTION**

**Method keywords** 

skin absorption
transport
kinetics
LC-MS
Scientific area keywords
pharmacokinetics
toxicokinetics
toxicity
Method description
Transport kinetics accross the skin and/or mucosa is investigated using Franz diffusion cells and LC-UV/MS detection of the investigated molecule in the receptor compartment.
Lab equipment
Franz diffusion cell system;
LC-UV/MS.
Method status
Internally validated
Published in peer reviewed journal
PROS, CONS & FUTURE POTENTIAL
Advantages

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

No in vitro cell line is used but real ex vivo skin or mucosa.

Taevernier L, et al. (2016). Human skin permeation of emerging mycotoxins (beauvericin and enniatins). J Expo Sci Environ Epidemiol. 26(3):277-87.

Taevernier L, et al. (2015). Enniatin-containing solutions for oromucosal use: Quality-by-design exvivo transmucosal risk assessment of composition variability. Int J Pharm. 491(1-2):144-51.

Veryser L, et al. (2014). Quantitative transdermal behavior of pellitorine from Anacyclus pyrethrum extract. Phytomedicine. 21(14):1801-7.

Veryser L, et al. (2016). Mucosal and blood-brain barrier transport kinetics of the plant N-alkylamide spilanthol using in vitro and in vivo models. BMC Complement Altern Med. 16:177.

Coordinated by









